

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 401/12	A1	(11) International Publication Number: WO 97/22603 (43) International Publication Date: 26 June 1997 (26.06.97)
(21) International Application Number: PCT/SE96/01603 (22) International Filing Date: 5 December 1996 (05.12.96) (30) Priority Data: 9504503-5 15 December 1995 (15.12.95) SE (71) Applicant (for all designated States except US): ASTRA AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): GUSTAVSSON, Anders [SE/SE]; Fruktvägen 52, S-155 31 Nykvam (SE). KÄLLSTRÖM, Åke [SE/SE]; Rönnbärsvägen 3, S-152 52 Södertälje (SE). (74) Agent: ASTRA AKTIEBOLAG; Patent Dept., S-151 85 Södertälje (SE).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: METHOD FOR THE SYNTHESIS OF A BENZIMIDAZOLE COMPOUND		
(57) Abstract A process for the manufacture of omeprazole from pyrmethyl alcohol via pyrmethyl chloride and pyrmetazole characterized in that the whole reaction sequence is carried out without any isolation or purification of intermediates. Further that the reaction is carried out in a main solvent system common for the whole reaction sequence and inert to the reactants formed during the process and used in the process. The process according to the present invention may also include an additional purification step.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

METHOD FOR THE SYNTHESIS OF A BENZIMIDAZOLE COMPOUND

Field of the invention

5 The present invention relates to a novel process for the synthesis of 5-methoxy-2-[[[4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl]sulphinyl]-1H-benzimidazole, known under the generic name omeprazole. Moreover, the present invention also relates to manufacture of a pharmaceutical preparation thereof and its use in medicine.

10 Background of the invention and prior art.

Omeprazole is an inhibitor of gastric acid secretion making it useful as an antiulcer agent. US 4 255 431 (corresponding to EP 0 005 129) discloses a process for the preparation of this class of substituted benzimidazoles. Said process comprises a couple of reaction steps.

15 The three last steps of the process utilize more than one solvent and requires isolation of intermediates to give the final product omeprazole. The resulting product is contaminated with starting materials and by-products.

WO 91/18895 describes an improved process for synthesis of omeprazole. This process
20 describes the oxidation step and a work-up procedure of omeprazole. The oxidation step utilizes m-chloroperoxybenzoic acid in a methylene chloride/water system at substantially constant pH of about 8.0 to 8.6. The work-up procedure includes crystallization of omeprazole by the addition of an alkyl formate.

25 Another process for the manufacture of omeprazole is described in US 5 391 752. This process also describes only the oxidation step. Said oxidation step utilizes magnesium monoperoxyphthalate as an oxidizing agent. The resulting product is contaminated with starting material and by-products.

Yet another process for the manufacture of compounds showing structural similarities to omeprazole is disclosed in US 4 619 997. The intermediates are isolated, but not purified, and different solvent systems are used in the chemical steps throughout the process.


- 5 In the light of the above there was still a need for a new convenient and more efficient process to manufacture omeprazole.

Summary of the invention.

- 10 In the following the improved process is described. Scheme I below describes three main reaction sequences in the manufacture of omeprazole. In the scheme the complete chemical names as well as short names for the starting material and the different intermediates are stated. The starting material as well as the reactants used are readily available by procedures described in the literature.

15

Step 1:



(4-Methoxy-3,5-dimethyl-
2-pyridinyl)methyl alcohol


(Pyrmethyl alcohol)

+

SOCl_2 (or another suitable chlorinating agent)

20


3



(4-Methoxy-3,5-dimethyl-
2-pyridinyl)methyl chloride


(Pyrmethyl chloride)

Step 2:



(4-Methoxy-3,5-dimethyl-
2-pyridinyl)methyl chloride

(Pyrmethyl chloride)




2-Mercapto-5-methoxybenzimidazole

(Metmercazole)

+

5




5-Methoxy-2-(((4-methoxy-3,5-
dimethyl-2-pyridinyl)-methyl)-
thio)-1H-benzimidazole

(Pyrmatazole)


Step 3:

10




5-Methoxy-2(((4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl)-thio)-1H-benzimidazole

(Pyrimetazole)



m-Chloroperoxybenzoic acid



5-Methoxy-2(((4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl)sulfinyl)-1H-benzimidazole

(Omeprazole)

Scheme I

Advantages of the improved process of the present invention are disclosed in the following paragraphs.

One object of the present invention is to provide a process for the manufacture of omeprazole with all three reaction steps, described above, being carried out without any isolation or purification of the intermediates, and carried out in one main solvent system common for the whole sequence. Such a process eliminates time consuming steps for isolation or purification of intermediates or solvent change in the process and thus making the process more efficient with a high production capacity. A further important benefit of

such a process is that the handling of toxicological and suspected mutagenic intermediates such as pyrmethyl chloride, due to isolation or purification, is eliminated.

Another preferred object of the present invention is to provide a process utilizing a solvent system which is environmentally friendly to avoid outlet of harmful solvents. There is a general interest from environmental groups both inside and outside the pharmaceutical industry that the industry shall develop and use environmentally friendly processes. In some countries the authorities also have made restrictions on the outlet of harmful chlorinated solvents into the air.

Surprisingly, the novel process of the present invention gives omeprazole in a high purity without any isolation or purification of intermediates. The omeprazole obtained may optionally be further processed in a purification step and/or optionally processed into pharmaceutically acceptable salts.

Another object of the present invention is to provide a process for an optional purification step of omeprazole.

Further, the present reaction sequence starting from pyrmethyl alcohol via pyrmethyl chloride and pyrmatazole giving omeprazole can be carried out in a main solvent system used through the whole reaction sequence. The solvent system is preferably inert against all reactants and suitable for the three process steps.

Detailed description of the invention.

In step 1 pyrmethyl alcohol is reacted with an excess of thionyl chloride or another suitable chlorinating agent giving alkyl chlorides, i.e. pyrmethyl chloride. The reaction is preferably carried out at ambient temperature and for approximately a period of 30 minutes. The reaction may also be performed at a temperature of -5°C to 25°C.

The reaction is carried out in a solvent system, which is used throughout the whole reaction sequence described in scheme I, i.e. from pyrmethyl alcohol to omeprazole. Such a solvent system suitable for the present process comprises a main solvent and optionally one or more co-solvents. The main solvent is preferably water immiscible, e.g. carbontetrachloride, 1,1,2-trichloroethane, chloroform, methylene chloride or toluene. Toluene is especially preferred from an environmental point of view.

Furthermore, the solvent system may optionally comprise co-solvent(s) to further improve/enhance the solubility of the reactants or products during the whole reaction sequence. The solvent system will function without any addition of a co-solvent, but the addition of such a co-solvent will enhance the process capacity, i.e. the yield per volume. Preferred co-solvents are alcohols and especially a lower alcohol comprising 1 - 4 carbon atoms, which carbon chain may be branched or straight.

Examples of solvent systems include, but are not limited to, 70-80% (by volume) methylene chloride or toluene and 20-30% (by volume) lower alcohol, e.g. ethanol.

In step 2, the pyrmethyl chloride formed in step 1 is reacted with metmercazole under alkaline conditions, e.g. an alkaline aqueous solution of metmercazole is prepared and mixed with the pyrmethyl chloride in the presence of a phase transfer catalyst. The reaction is preferably carried out at a temperature of 30 - 60°C during a prolonged period of time.

Metmercazole is charged in approximately stoichiometric amount to the pyrmethyl chloride. A phase transfer catalyst may be used, for instance a tertiary amine, preferably tetrabutylammonium bromide.

The two phases formed are separated and the aqueous phase may be extracted with the used water immiscible organic solvent, such as methylene chloride or toluene. Furthermore, an alcohol may be added to the pyrmatazole solution to improve the solubility of the formed intermediate, pyrmatazole.

In step 3, m-chloroperoxybenzoic acid is dissolved in an water immiscible organic solvent such as methylene chloride or toluene. The m-chloroperoxybenzoic acid is charged to the obtained pyrimetazole in a buffered two phase system, consisting of the water immiscible solvent, such as methylene chloride or toluene, and aqueous base, such as sodium or potassium hydrogencarbonate. The organic solvent may optionally be diluted with an alcohol to enhance the solubility of the m-chloroperoxybenzoic acid. The oxidation with m-chloroperoxybenzoic acid is preferably performed between 0° and 25°C. The obtained product is extracted into an alkaline, aqueous phase and the formed product is precipitated from the aqueous phase by lowering the pH of the solution to approximately 9 at ambient temperature. Omeprazole is isolated and washed. The overall yield in the process is usually higher than 75%.

The additional purification can be carried out by crystallization from one or several organic solvents, or by precipitation from alkaline water by lowering the pH through addition of an acid or an alkyl formate.

Examples of suitable acids include, but are not limited to, carboxylic acids such as formic acid, acetic acid, or citric acid, and mineral acids, such as hydrogen chloride, hydrogen bromide, hydrogen fluoride, hydrogen iodide, sulfuric acid, nitric acid, or phosphoric acid. Preferred acids are carboxylic acids. Suitable alkyl formates include, but are not limited to, methyl formate and ethyl formate. A preferred alkyl formate is methyl formate.

The present invention is described more in detail in the following non-limiting examples.

Manufacturing of omeprazole.

Example 1.

17.8 g (0.15 mol) Thionyl chloride in 13 mL methylene chloride was charged to a methylene chloride solution of 16.8 g (0.10 mol) pyrmethyl alcohol at ambient temperature. The solution was stirred for approximately 30 minutes.

5 18.0 g (0.10 mol) Metmercazone, 47.3 g (0.58 mol) aqueous sodium hydroxide (50% w/w) and tetrabutylammonium bromide (0.9 g) were mixed at ambient temperature. The pyrmethyl chloride solution was added at a temperature between 25° - 40°C and the mixture was refluxed for 1 - 2 hours. The two phases were separated and the aqueous phase was washed with methylene chloride. The organic phases were combined.

10

10.4 g (0.10 mol) Potassium bicarbonate dissolved in water was added to the pyrmatazone solution.

22.3 g (0.099 mol) *Meta*-chloroperoxybenzoic acid (76.5% w/w) dissolved in 42 mL of
15 methylene chloride and 10 mL of ethanol were charged to the prepared mixture of pyrmatazone in methylene chloride and potassium bicarbonate in water. The oxidation was carried out at 0 - 15°C. The product was extracted into the aqueous phase at an alkaline pH by addition of sodium hydroxide. The two phases were separated and the organic phase was washed with an alkaline water solution. The aqueous phases were combined.

20

Omeprazole was precipitated from the aqueous phase by the addition of 12 mL methyl formate, filtered and washed with water/methanol and dried. Omeprazole was obtained in a yield of 76%.

25 Example 2.

13.1 g (0.11 mol) Thionyl chloride was charged to a toluene solution of 16.7 g (0.10 mol) pyrmethyl alcohol at ambient temperature. The reaction was continued for approximately 30 minutes.

30

The formed pyrmethyl chloride in toluene was mixed with an alkaline aqueous solution of 18.0 g (0.10 mol) metmercazole. The reaction was performed in the presence of 0.9 g tetrabutylammonium bromide at 40°C during a couple of hours. The two phases were separated and the aqueous phase was extracted with toluene. Ethanol was added to improve
5 the solubility of the resulting product before the solution was cooled to ambient temperature.

24.4 g (0.10 mol) *Meta*-chloroperoxybenzoic acid in 42 mL toluene and 10 mL ethanol were charged to a two phase system prepared from the obtained pyrmetazole toluene
10 solution, and potassium bicarbonate (13.8 g) in water (120 mL). The oxidation was performed between 0°C and room temperature. The product was extracted into an alkaline aqueous phase by the addition of sodium hydroxide. The two phases were separated and the organic phase was washed with water. Omeprazole was precipitated by lowering the pH of the solution to approximately 9 at ambient temperature. Omeprazole was isolated, washed,
15 and dried. The overall yield was approximately 77%.

Additional purification of omeprazole.

20 Example 3

20.0 g (0.0579 moles) of omeprazole, prepared according to example 1, 125 mL of methanol and 54 mL of water were mixed at ambient temperature.

25 The product was dissolved at an alkaline pH by addition of sodium hydroxide. The solution was filtered and cooled to approximately 0°C.

The product was precipitated by addition of diluted acetic acid, filtered and washed with water/methanol, and thereafter dried. Omeprazole was obtained in a yield of 90 %.

Example 4

11.2 g (0.0324 moles) of omeprazole, prepared according to example 1, 69 mL of
5 methanol, 34 mL of methylene chloride and 0.3 mL of ammonia were mixed at ambient
temperature and the obtained solution was filtered.

The product was precipitated by evaporation of methylene chloride under reduced pressure.

10 0.3 mL of ammonia was added and the slurry was cooled to 0°C.

The product was filtered, washed with methanol containing approximately 1% v/v of
ammonia, and dried. Omeprazole was obtained in a yield of 90 %.


15

With respect to the environmental point of view, the best way to carry out one aspect of the
present invention is according to the process described in Example 2.

CLAIMS.

1. A process for the manufacture of 5-methoxy-2-[[(4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl]sulphinyl]-1H-benzimidazole from (4-methoxy-3,5-dimethyl-2-pyridinyl)-
5 methyl alcohol comprising the following reaction steps

Step 1:




(4-Methoxy-3,5-dimethyl-
2-pyridinyl)methyl alcohol

(Pyrmethyl alcohol)

+

SOCl_2 (or another suitable chlorinating agent)


10



(4-Methoxy-3,5-dimethyl-
2-pyridinyl)methyl chloride


(Pyrmethyl chloride)

Step 2:



(4-Methoxy-3,5-dimethyl-2-pyridinyl)methyl chloride

(Pyrmethyl chloride)




2-Mercapto-5-methoxybenzimidazole

(Metmercazole)

+


5



5-Methoxy-2(((4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl)-thio)-1H-benzimidazole

(Pyrmatazole)


Step 3:



5-Methoxy-2(((4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl)-thio)-1H-benzimidazole


(Pyrmatazole)

+



m-Chloroperoxybenzoic acid

10



5-Methoxy-2-(((4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl)sulfinyl)-1H-benzimidazole

(Omeprazole)

characterized in that the reaction steps are carried out in a consecutive order without isolation of the intermediates formed during the process using one main solvent system
5 common for the whole reaction sequence.

2. A process according to claim 1, characterized in that the solvent system comprises a main solvent and optionally one or more co-solvents.

10

3. A process according to claim 2, characterized in that the main solvent is methylene chloride.

4. A process according to claim 2, characterized in that the main solvent is toluene.

15

5. A process according to claim 2, characterized in that the co-solvent is an alcohol.

6. A process according to claim 1, characterized in that step 1 is carried out at a temperature between -5°C to 25°C.

20

7. A process according to claim 1, characterized in that step 2 is carried out with a phase transfer catalyst.

8. A process according to claim 7, characterized in that the phase transfer catalyst is tetrabutylammonium bromide.

9. A process according to claim 1, characterized in that step 2 is carried out at a
5 temperature between 30°C to 60°C.

10. A process according to anyone of the proceeding claims wherein the resulting 5-methoxy-2-[[4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl]sulphinyl]-1H-benzimidazole is further purified in a purification step.

10

11. A process according to claim 10, characterized in that the purification step involves crystallization from one or several organic solvents.

12. A process according to claim 10, characterized in that the purification step involves
15 precipitation from alkaline water by lowering the pH through addition of an acid or an alkyl formate.

13. A pharmaceutical preparation comprising 5-methoxy-2-[[4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl]sulphinyl]-1H-benzimidazole and a pharmaceutically acceptable carrier
20 or diluent characterized in that the 5-methoxy-2-[[4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl]sulphinyl]-1H-benzimidazole is prepared by a process according to one of the claims 1-12.

14. 5-Methoxy-2-[[4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl]sulphinyl]-1H-benzimidazole prepared by a process according to one of claims 1-12.

1
INTERNATIONAL SEARCH REPORTInternational application No.
PCT/SE 96/01603

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07D 401/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4619997 A (JOHN C. SIH), 28 October 1986 (28.10.86), see column 9, line 9 - line 17 and column 7 examples 1 and 2 ---	1-12
A	US 4255431 A (ULF K. JUNGREN ET AL), 10 March 1981 (10.03.81), see example 22 and method C	1-12
X	-----	13, 14

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

- * Special categories of cited documents:
- "A" document defining the general state of the art which is not considered to be of particular relevance
 - "E" earlier document but published on or after the international filing date
 - "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 - "O" document referring to an oral disclosure, use, exhibition or other means
 - "P" document published prior to the international filing date but later than the priority date claimed
 - "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 - "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 - "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
 - "&" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

30 January 1997

01-03-1997

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Göran Karlsson
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

Information on patent family members

28/10/96

International application No.

PCT/SE 96/01603

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 4619997	28/10/86	AU-B- 571907	28/04/88
		AU-A- 3564384	13/06/85
		EP-A,B- 0150586	07/08/85
		SE-T3- 0150586	
		IE-B- 58431	22/09/93
		US-A- 4575554	11/03/86
US-A- 4255431	10/03/81	AT-B- 374471	25/04/84
		AT-B- 374472	25/04/84
		AT-B- 374473	25/04/84
		AT-B- 375365	25/07/84
		AT-B- 389995	26/02/90
		AU-B- 529654	16/06/83
		AU-A- 4602779	18/10/79
		CA-A- 1127158	06/07/82
		CA-A- 1129417	10/08/82
		CY-A- 1232	29/06/84
		EP-A,B- 0005129	31/10/79
		SE-T3- 0005129	
		HK-A- 15284	02/03/84
		JP-C- 1312930	28/04/86
		JP-C- 1504537	13/07/89
		JP-A- 54141783	05/11/79
		JP-A- 58192880	10/11/83
		JP-B- 60034956	12/08/85
		JP-B- 63053191	21/10/88
		LU-A- 88305	04/05/94
		LU-A- 88307	04/05/94
		SE-A- 7804231	15/10/79
		SU-A- 895292	30/12/81
		US-A- 4337257	29/06/82
		US-A- 4508905	02/04/85